



アナストロゾール+ラパチニブ療法

1日1回 (HER2陽性)

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
アナストロゾール (1mg) 1日1回経口(連日投与)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓
ラパチニブ (1,500mg) 1日1回経口(連日投与) [※]	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓

※食事の1時間以上前もしくは食後1時間以降に経口投与する

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
血栓塞栓症																												
肝機能障害																												
関節痛																												
性器出血																												
高コレステロール血症																												
ほてり																												
頭痛																												
嘔気																												
発疹																												
骨粗しょう症																												
下痢																												
脱毛症																												
皮膚乾燥																												
爪の障害																												
疲労																												

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)