



アナストロゾール+パルボシクリブ療法 4週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	
アナストロゾール (1mg) 1日1回経口(連日投与)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	
パルボシクリブ (125mg) 1日1回経口※ (3週連続/1週休薬)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓										↓	↓	↓	↓

※食後投与

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	
血栓塞栓症																																	
肝機能障害																																	
関節痛																																	
性器出血																																	
高コレステロール血症																																	
ほてり																																	
頭痛																																	
嘔気																																	
発疹																																	
骨粗しょう症																																	
骨髄抑制																																	
間質性肺疾患																																	
脱毛																																	
疲労																																	
口内炎																																	
下痢																																	

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)