



ビカルタミド+リュープロレリン療法

4週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	
ビカルタミド (80mg) 1日1回経口(連日投与)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	
リュープロレリン (3.75mg) 皮下注	↓																																↓

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32		
肝機能障害																																		
白血球減少																																		
血小板減少																																		
乳房腫脹																																		
乳房圧痛																																		
ほてり																																		
発汗																																		
勃起力低下																																		
うつ状態																																		
骨疼痛																																		
脊髄圧迫																																		
熱感																																		
硬結(投与部位)																																		

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)