



ビカルタミド+デガレリクス療法 4週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32
ビカルタミド (80mg) 1日1回経口(連日投与)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓
デガレリクス (初回240mg、2回目以降80mg) 皮下注	↓																											↓				

※初回は240mgを1カ所あたり120mgずつ腹部2カ所に皮下投与する。2回目以降は初回投与4週間後より維持用量を投与する。4週間間隔で投与する場合は80mgを腹部1カ所に皮下投与する。

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32																																
肝機能障害																																																																
白血球減少																																																																
血小板減少																																																																
乳房腫脹																																																																
乳房圧痛																																																																
ほてり																																																																
発汗																																																																
勃起力低下																																																																
注射部位疼痛																																																																
体重増加																																																																
発熱																																																																
高血圧																																																																

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)