



## レジメン&副作用情報

ホリナート>大腸がん

### UFT+ホリナート療法

28日間投与、7日間休薬（術後補助化学療法の場合は5コース）

レジメン/(日)	1	8	15	22	29	36	43	50	57	64	71	78	85	92	99
UFT (1回100mg/m <sup>2</sup> ) 1日3回経口	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓
ホリナート (1回25mg) 1日3回経口	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓

副作用/(日)	1	8	15	22	29	36	43	50	57	64	71	78	85	92	99	
骨髄抑制		[Green shaded area]					[Green shaded area]					[Green shaded area]				
黄だん・肝障害		[Green shaded area]					[Green shaded area]					[Green shaded area]				
下痢		[Green shaded area]					[Green shaded area]					[Green shaded area]				
嘔気・嘔吐	[Green shaded area]					[Green shaded area]					[Green shaded area]					
口内炎		[Green shaded area]					[Green shaded area]					[Green shaded area]				
食欲不振	[Green shaded area]					[Green shaded area]					[Green shaded area]					
嗅覚障害・味覚異常		[Green shaded area]					[Green shaded area]					[Green shaded area]				

\* UFT(テガフル・ウラシル)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)