



レトロゾール+ラパチニブ療法 1日1回 (HER2陽性)

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
レトロゾール (2.5mg) 1日1回経口(連日投与)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓
ラパチニブ (1,500mg) 1日1回経口(連日投与)*	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓

\*食事の1時間以上前もしくは食後1時間以降に経口投与する

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
血栓症																													
狭心症																													
肝機能障害																													
コレステロール値増加																													
ほてり																													
頭痛																													
関節痛																													
悪心																													
発疹																													
掻痒症																													
骨粗しょう症																													
下痢																													
脱毛症																													
皮膚乾燥																													
爪の障害																													
疲労																													

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)