



### レトロゾール+アベマシクリブ療法

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
レトロゾール (2.5mg) 1日1回経口(連日投与)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓
アベマシクリブ (1回 150mg) 1日2回経口(連日投与)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
血栓症																													
狭心症																													
肝機能障害																													
コレステロール値増加																													
ほてり																													
頭痛																													
関節痛																													
悪心																													
発疹																													
掻痒症																													
骨粗しょう症																													
下痢																													
骨髄抑制																													
疲労																													
腹痛																													
嘔吐																													
脱毛																													
食欲減退																													

注)アベマシクリブの重大な副作用として、間質性肺疾患があらわれることがあるので、観察を十分に行い、異常が認められた場合には、本剤を中止し、必要に応じて、胸部CT、血清マーカー等の検査を実施するとともに、適切な処置を行うこと。

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)