



mFOLFOX6+ベバシズマブ療法 2週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
ベバシズマブ (5mg/kg) 点滴静注(初回90分)*	↓														↓													
オキサリプラチン (85mg/m ²) 点滴静注(120分)	↓														↓													
レボホリナート (200mg/m ²) 点滴静注(120分)	↓														↓													
5-FU (400mg/m ²) 急速静注(5分)	↓														↓													
5-FU (2,400mg/m ²) 持続静注(46時間)	→														→													

※初回投与で時間90分、2回目60分、3回目以降は30分への短縮も可能

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
嘔気・嘔吐	■														■													
食欲不振	■														■													
過敏症	■														■													
骨髄抑制															■													
口内炎			■															■										
手足症候群		■																										
末梢神経障害		■																										
消化管穿孔	■																											
創傷治癒遅延	■																											
出血・塞栓症	■																											
高血圧脳症	■																											

* 5-FU(フルオロウラシル)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)