



FOLFIRI療法

2週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
イリノテカン (150mg/m ²) 点滴静注(90分)	↓														↓													
レボホリナート (200mg/m ²) 点滴静注(120分)	↓														↓													
5-FU (400mg/m ²) 急速静注(5分)	↓														↓													
5-FU (2,400mg/m ²) 持続静注(46時間)	→														→													

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
嘔気・嘔吐	[Green wedge]														[Green wedge]																
食欲不振	[Green wedge]														[Green wedge]																
過敏症	[Green semi-circle]														[Green semi-circle]																
骨髄抑制										[Green oval]												[Green oval]									
口内炎		[Green oval]														[Green oval]															
手足症候群			[Green oval]																												
脱毛			[Green bar]																												
下痢		[Green oval]														[Green oval]															

* 5-FU(フルオロウラシル)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)