



FOLFIRI+セツキシマブ療法 (KRAS遺伝子野生型)

2週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
セツキシマブ (400→250mg/m ²)※ 点滴静注(120→60分)	↓							↓							↓							↓							
イリノテカン (150mg/m ²) 点滴静注(90分)	↓														↓														
レボホリナート (200mg/m ²) 点滴静注(120分)	↓														↓														
5-FU (400mg/m ²) 急速静注(5分)	↓														↓														
5-FU (2,400mg/m ²) 持続静注(46時間)	→														→														

※初回投与のみ400mg/m² 点滴静注120分、2回目以降は250mg/m² 点滴静注60分

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28		
嘔気・嘔吐	■														■															
食欲不振	■														■															
過敏症	■														■															
骨髄抑制										■													■							
口内炎		■																■												
手足症候群			■																											
脱毛															■															
下痢		■																■												
ざ瘡	■																													
皮膚乾燥	■																													
発疹	■																													
痒痒症	■																													

*5-FU(フルオロウラシル)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)