



FOLFIRI+ラムシルマブ療法

2週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
ラムシルマブ (8mg/kg) 点滴静注(60分)	↓														↓													
イリノテカン (150mg/m ²) 点滴静注(90分)	↓														↓													
レボホリナート (200mg/m ²) 点滴静注(120分)	↓														↓													
5-FU (400mg/m ²) 急速静注(5分)	↓														↓													
5-FU (2,400mg/m ²) 持続静注(46時間)	→														→													

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28				
嘔気・嘔吐	■														■																	
食欲不振	■														■																	
過敏症	■														■																	
骨髄抑制											■											■										
口内炎		■																■														
手足症候群			■																													
脱毛															■																	
下痢		■																■														
出血・塞栓症			■																													
消化管穿孔	■																															
創傷治癒障害	■																															

* 5-FU(フルオロウラシル)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)