



FOLFIRI+パニツムマブ療法 (KRAS遺伝子野生型)

2週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
パニツムマブ (6mg/kg) 点滴静注(60分)	↓														↓														
イリノテカン (150mg/m ²) 点滴静注(90分)	↓														↓														
レボホリナート (200mg/m ²) 点滴静注(120分)	↓														↓														
5-FU (400mg/m ²) 急速静注(5分)	↓														↓														
5-FU (2,400mg/m ²) 持続静注(46時間)	→														→														

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28			
嘔気・嘔吐	■														■																
食欲不振	■														■																
過敏症	■														■																
骨髄抑制										■													■								
口内炎			■																■												
手足症候群					■																										
脱毛															■																
下痢			■																■												
ざ瘡様皮膚炎	■														■																
爪囲炎															■																
皮膚乾燥															■																
掻痒	■																														

* 5-FU(フルオロウラシル)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)

Empty rounded rectangular box for contact information.