



XELOX療法

3週間ごと（術後補助化学療法の場合は8コース）

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
オキサリプラチン (130mg/m ²) 点滴静注(120分)	↓																					↓							
カペシタビン (1回1,000mg/m ²)※ 1日2回経口	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓								↓	↓	↓	↓	↓	↓	↓	

※カペシタビン投与量: 体表面積 1.36m²未満1回1,200mg、1.36m²以上1.66m²未満1回1,500mg、1.66m²以上1.96m²未満1回1,800mg、1.96m²以上1回2,100mg

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28								
骨髄抑制										[Green shaded area]																										
疲労	[Green shaded area]																																			
腹痛	[Green shaded area]																																			
下痢										[Green shaded area]																										
嘔気・嘔吐	[Green shaded area]																																			
口内炎				[Green shaded area]																																
食欲不振	[Green shaded area]																																			
末梢神経障害																																				
発疹			[Green shaded area]																																	
手足症候群			[Green shaded area]																																	

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)
