



SOX療法

3週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
S-1 (1回40mg) [※] 1日2回経口(14日間)	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓								↓	↓	↓	↓	↓	↓	↓
オキサリプラチン (130mg/m ²) 点滴静注(120分)	↓																					↓						

※S-1初回基準投与量:体表面積1.25m²未満 40mg/回、1.25m²以上1.5m²未満 50mg/回、1.5m²以上 60mg/回

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
嘔気・嘔吐	[Green shaded area]																					[Green shaded area]							
食欲不振	[Green shaded area]																						[Green shaded area]						
下痢	[Green shaded area]																												
骨髄抑制																													
口内炎																													
皮疹																													
肝機能障害																													
末梢神経障害																													

* S-1(テガフル・ギメラシル・オテラシルカリウム配合剤)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)