



mFOLFOX6+パニツムマブ療法 (KRAS遺伝子野生型)

2週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
パニツムマブ (6mg/kg) 点滴静注(60分)	↓														↓													
オキサリプラチン (85mg/m ²) 点滴静注(120分)	↓														↓													
レボホリナート (200mg/m ²) 点滴静注(120分)	↓														↓													
5-FU (400mg/m ²) 急速静注(5分)	↓														↓													
5-FU (2,400mg/m ²) 持続静注(46時間)	→														→													

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28											
嘔気・嘔吐	■														■																								
食欲不振	■														■																								
過敏症	■														■																								
骨髄抑制													■																										
口内炎			■															■																					
手足症候群			■																																				
末梢神経障害			■																																				
ざ瘡様皮膚炎	■														■																								
爪囲炎															■																								
皮膚乾燥															■																								
掻痒	■																																						

* 5-FU(フルオロウラシル)

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)