



XELOX + ベバシズマブ療法

3週間ごと

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
ベバシズマブ (7.5mg/kg) 点滴静注(初回90分) ^{※1}	↓																					↓						
オキサリプラチン (130mg/m ²) 点滴静注(120分)	↓																					↓						
カペシタビン (1回1,000mg/m ²) ^{※2} 1日2回経口	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓								↓	↓	↓	↓	↓	↓	↓

※1 初回投与時間90分、2回目60分、3回目以降は30分への短縮も可能

※2 カペシタビン投与量: 体表面積 1.36m²未満1回1,200mg、1.36m²以上1.66m²未満1回1,500mg、1.66m²以上1.96m²未満1回1,800mg、1.96m²以上1回2,100mg

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
骨髄抑制																													
疲労																													
腹痛																													
下痢																													
嘔気・嘔吐																													
口内炎																													
食欲不振																													
末梢神経障害																													
発疹																													
手足症候群																													
消化管穿孔																													
創傷治癒遅延																													
出血・塞栓症																													
高血圧脳症																													

※副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)