



Tamoxifen単独療法

1日1回（術後内分泌療法の場合は5年～10年）

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
Tamoxifen (20mg) 1日1回経口(連日投与) [※]	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓

※1日最高量は40mgまで。錠10mg投与時は1日20mgを1～2回に分割経口投与する。

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
血栓塞栓症																													
肝機能異常																													
ほてり・潮紅																													
発汗																													
無月経・月経異常																													
性器出血																													
膣分泌物																													
悪心・嘔吐																													
食欲不振																													
頭痛																													
めまい																													
発疹																													
体重増加																													

* Tamoxifen(タモキシフェン)

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)