



Tamoxifen+Leuprorelin療法 4週間ごと（閉経前乳がん）

レジメン/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	
Tamoxifen (20mg) 1日1回経口(連日投与)*	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	↓	
Leuprorelin (3.75 mg) 皮下注	↓																												↓				

※1日最高量は40mgまで。錠10mg投与時は1日20mgを1~2回に分割経口投与する。

副作用/(日)	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	
血栓塞栓症																																	
肝機能異常																																	
ほてり・潮紅																																	
発汗																																	
無月経・月経異常																																	
性器出血																																	
膣分泌物																																	
悪心・嘔吐																																	
食欲不振																																	
頭痛																																	
めまい																																	
発疹																																	
体重増加																																	
更年期障害様のうつ状態																																	
肩こり																																	
不眠																																	
関節痛																																	
骨疼痛																																	
硬結(投与部位)																																	

* Tamoxifen(タモキシフェン)、Leuprorelin(リュープロレリン)

副作用は、あくまで一般的な目安であり、実際の発現頻度/程度、時期については個人差があります。

連絡先(医療機関名)

Blank box for entering the medical institution name.